

WO9411384

Publication Title:

NOVEL CYTOKINE DESIGNATED ELK LIGAND

Abstract:

Elk ligand (elk-L) polypeptides as well as DNA sequences, vectors and transformed host cells useful in providing elk-L polypeptides. The elk-L polypeptide binds to a cell surface receptor that is a member of the tyrosine kinase receptor family.

Data supplied from the esp@cenet database - <http://ep.espacenet.com>

This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/04, C12N 15/12 C07K 13/00, 15/28	A1	(11) International Publication Number: WO 94/11384 (43) International Publication Date: 26 May 1994 (26.05.94)
(21) International Application Number: PCT/US93/10955 (22) International Filing Date: 15 November 1993 (15.11.93) (30) Priority data: 07/977,693 13 November 1992 (13.11.92) US (71) Applicant: IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US). (72) Inventors: LYMAN, Stewart ; 312 North 45th Street, Seat- tle, WA 98103 (US). BECKMANN, M., Patricia ; 15875 Nesika Bay Road, Poulsbo, WA 98370 (US). BAUM, Pe- ter, R. ; 6027 - 34th Avenue N.E., Seattle, WA 98115 (US).		(74) Agent: SEESE, Kathryn, A.; Immunex Corporation, 51 University Street, Seattle, WA 98101 (US). (81) Designated States: AU, BR, CA, JP, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: NOVEL CYTOKINE DESIGNATED ELK LIGAND (57) Abstract Elk ligand (elk-L) polypeptides as well as DNA sequences, vectors and transformed host cells useful in providing elk-L polypeptides. The elk-L polypeptide binds to a cell surface receptor that is a member of the tyrosine kinase receptor family.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

TITLE**NOVEL CYTOKINE DESIGNATED ELK LIGAND**

10

BACKGROUND OF THE INVENTION

The cell surface protein designated elk is a member of a family of proteins known as the tyrosine kinase receptors. Proteins of this family have an intrinsic kinase activity that is activated upon ligand binding (Ullrich et al., *Cold Spring Harbor Symp. Quant. Biol.* 51:713, 1986). A partial clone of elk was first discovered in a rat brain cDNA expression library that was screened for proteins expressing tyrosine kinase activity (Letwin et al., *Oncogene* 3:621, 1988). Later, a composite sequence spanning the entire elk coding region was derived from partial clones isolated from a rat brain cDNA library and a rat cerebellar brain library using the partial clone as a probe (Lhotak et al., *Mol. Cell. Biol.* 11:2496, 1991). The elk protein is very closely related to a number of other receptor tyrosine kinases, including hek (Boyd et al. *J. Biol. Chem.* 267:3262, 1992 and Wicks et al. *Proc. Natl. Acad. Sci. USA* 89:1611, 1992); the hek homologs mek4 and cek4 (Sajjadi et al. *New Biol.* 3:769, 1991); eek (Chan et al. *Oncogene* 6:1057, 1991); erk (Chan et al. *supra.*), eck (Lindberg et al. *Mol. Cell. Biol.* 10:6316, 1990); cek5 (Pasquale, E.B. *Cell Regulation* 2:523, 1991); and eph (Hirai et al. *Science* 238:1717, 1987). The proteins of this subfamily are related not only in their cytoplasmic domains, but also in their extracellular domains, which are 41 to 68% identical. Interestingly, the tissue distributions of these various receptors are diverse. For example, expression of elk mRNA has been shown to be limited to testis and brain (Lhotak et al., *supra.*), whereas eck is found not only in these same two tissues but in lung, intestine, kidney, spleen, ovary, and skin as well.

Ligands for the receptor tyrosine kinases are a diverse group of proteins that affect the growth, differentiation, and survival of cells expressing the receptors. To date, no ligand for elk has been discovered. Identification of the putative ligand would prove useful in investigating the nature of cellular processes that might be regulated by the elk protein.

35

SUMMARY OF THE INVENTION

The present invention provides a novel cytokine designated elk ligand (elk-L) that binds to the rat cell surface receptor known as elk. The present invention also provides isolated DNA encoding the elk-L protein, expression vectors comprising the isolated DNA, and a method for producing elk-L by cultivating host cells containing the
5 expression vectors under conditions appropriate for expression of the elk-L protein. Antibodies directed against the elk-L protein or an immunogenic fragment thereof are also disclosed.

DETAILED DESCRIPTION OF THE INVENTION

10 A cDNA encoding a novel protein ligand that binds to the rat cell surface protein known as elk has been isolated in accordance with the present invention. Also provided are expression vectors comprising the elk ligand (elk-L) cDNA and methods for producing recombinant elk-L polypeptides by cultivating host cells containing the
15 expression vectors under conditions appropriate for expression of elk-L, and recovering the expressed elk-L. Purified elk-L protein is also encompassed by the present invention, including soluble forms of the protein comprising the extracellular domain.

The present invention also provides elk-L or antigenic fragments thereof that
20 can act as immunogens to generate antibodies specific to the elk-L immunogens. Monoclonal antibodies specific for elk-L or antigenic fragments thereof thus can be prepared.

The novel cytokine disclosed herein is a ligand for elk, a rat cell surface receptor that is a member of the tyrosine kinase receptor family. Members of this receptor
25 family have a kinase activity that is activated upon ligand binding. Binding of elk-L to elk on the cell surface may initiate a biological signal mediated by elk. One use of the elk ligand of the present invention is as a research tool for studying the nature of this biological signal and the role that elk-L, in conjunction with elk, may play in growth or differentiation of cells bearing the elk receptor. Expression of elk mRNA has been
30 detected in the brain and testis of rats (Lhotok et al., *supra*), and the possibility that elk is capable of oncogenic activation has been suggested (Letwin et al., *supra*).

The elk-L polypeptides of the present invention find use as protein purification reagents and also may be employed in *in vitro* assays for detection of elk or elk-L or the interactions thereof. These and other uses of elk ligands are further discussed below.

35 To identify cells suitable for use as nucleic acid sources in the cloning attempt, over 30 different types of murine and human cells were screened for the ability to bind elk (in the form of a fusion protein comprising rat elk and an antibody Fc polypeptide).

As described in Example 2, none of the cell types exhibited detectable elk binding. Since placental tissue is rich in growth and differentiation factors, a human placental cDNA expression library was screened with rat elk/Fc in an attempt to isolate an elk-L clone. Although it was not known whether or not placenta expressed an elk-L, and the ability of rat elk to bind to human elk-L also was unknown, human elk-L cDNA was successfully isolated as described in Example 3. The DNA sequence and encoded amino acid sequence of the coding region of a human elk-L cDNA clone are set forth in SEQ ID NO:1 and SEQ ID NO:2.

Human elk-L cDNA comprising the coding region was isolated from the positive clone and inserted into the Sma I site (in the multiple cloning site region) of cloning vector pBLUESCRIPT® SK(-), available from Stratagene Cloning Systems, La Jolla, CA. The resulting recombinant vector, designated tele 7 in pBLUESCRIPT® SK(-), in *E. coli* DH5α cells, was deposited with the American Type Culture Collection on October 9, 1992, and assigned accession no. ATCC 69085. The deposit was made under the terms of the Budapest Treaty.

Comparison of both the nucleotide and encoded amino acid sequences of the human elk-L cDNA clone with the Genbank and Swisspro databases showed that the sequence of the elk ligand was unique. One amino acid sequence was identified in this search that did share limited sequence identity with the elk ligand. That sequence was for the B61 protein, which has previously been identified as the product of a novel immediate-early response gene induced by TNF in human umbilical vein endothelial cells (Holzman et al., *Mol. Cell. Biol.* 10:5830, 1990). All four of the cysteine residues in the extracellular domain of the two proteins are conserved, and the overall amino acid identity between human elk-L and B61 is 33%. In contrast to the elk ligand, the B61 protein has been reported to be secreted, but terminates with a hydrophobic tail and has been suggested to be associated with the membrane through a glycosylphosphatidyl inositol linkage (Holzman et al., *supra*). The function of the B61 protein is unknown.

The term "elk-L" as used herein refers to a genus of polypeptides which are capable of binding elk and exhibit homology to the native elk-L polypeptides disclosed herein, preferably comprising an amino acid sequence that is over 80% identical to the amino acid sequence of SEQ ID NO:2. Human elk-L is within the scope of the present invention, as are elk-L proteins derived from other mammalian species including but not limited to murine, rat, bovine, porcine, or various primate cells. As used herein, the term "elk-L" includes membrane-bound proteins (comprising a cytoplasmic domain, a transmembrane region, and an extracellular domain) as well as truncated proteins that

retain the elk-binding property. Such truncated proteins include, for example, soluble elk-L comprising only the extracellular (receptor binding) domain.

The human elk-L cDNA may be radiolabeled and used as a probe to isolate other mammalian elk-L cDNAs by cross-species hybridization. For example, a cDNA library prepared from placental tissue of other mammalian species may be screened with radiolabeled human elk-L cDNA to isolate a positive clone. Alternatively, mRNAs isolated from various cell lines can be screened by Northern hybridization to determine a suitable source of mammalian elk-L mRNA for use in cloning an elk-L gene.

Although an elk/Fc fusion protein was employed in the screening procedure described in Example 3 below, elk can be used to screen clones and candidate cell lines for expression of elk-L proteins. The elk/Fc fusion protein, however, offers the advantage of being easily purified. In addition, disulfide bonds form between the Fc regions of two separate fusion protein chains, creating dimers. The dimeric elk/Fc receptor was chosen for the potential advantage of higher affinity binding of the elk ligand, in view of the possibility that the ligand being sought would be multimeric.

Other antibody Fc regions may be substituted for the human IgG1 Fc region described in Example 1. Other suitable Fc regions are those that can bind with high affinity to protein A or protein G, and include the Fc region of murine IgG1 or fragments of the human IgG1 Fc region, e.g., fragments comprising at least the hinge region so that interchain disulfide bonds will form.

One embodiment of the present invention provides soluble elk-L polypeptides. Soluble elk-L polypeptides comprise all or part of the extracellular domain of a native elk-L but lack the transmembrane region that would cause retention of the polypeptide on a cell membrane. Soluble elk-L polypeptides advantageously comprise the native (or a heterologous) signal peptide when initially synthesized to promote secretion, but the signal peptide is cleaved upon secretion of elk-L from the cell. The soluble elk-L polypeptides that may be employed retain the ability to bind the elk receptor. Soluble elk-L may also include part of the transmembrane region or part of the cytoplasmic domain or other sequences, provided that the soluble elk-L protein is capable of being secreted.

Soluble elk-L may be identified (and distinguished from its non-soluble membrane-bound counterparts) by separating intact cells which express the desired protein from the culture medium, e.g., by centrifugation, and assaying the medium (supernatant) for the presence of the desired protein. The presence of elk-L in the medium indicates that the protein was secreted from the cells and thus is a soluble form of the desired protein. Soluble elk-L may be a naturally-occurring form of this protein.

The use of soluble forms of elk-L is advantageous for certain applications. Purification of the proteins from recombinant host cells is facilitated, since the soluble proteins are secreted from the cells. Further, soluble proteins are generally more suitable for intravenous administration.

5 Examples of soluble elk-L polypeptides include those comprising the entire extracellular domain of a native elk-L protein. One such soluble elk-L protein comprises amino acids 1 (Ala) through 213 (Lys) of SEQ ID NO:2. When initially expressed within a host cell, the soluble protein may additionally comprise one of the heterologous signal peptides described below that is functional within the host cells
10 employed. Alternatively, the protein may comprise the native signal peptide, such that the elk-L comprises amino acids -24 (Met) through 213 (Lys) of SEQ ID NO:2. In one embodiment of the invention, soluble elk-L is initially expressed as a fusion protein comprising (from N- to C-terminus) the yeast α factor signal peptide, the FLAG[®] peptide described below and in U. S. Patent 5,011,912, and soluble elk-L comprising
15 amino acids 1-213 of SEQ ID NO:2. This recombinant fusion protein is expressed in and secreted from yeast cells. The FLAG[®] peptide facilitates purification of the protein, and subsequently may be cleaved from the soluble elk-L using bovine mucosal enterokinase. DNA sequences encoding soluble elk-L proteins are encompassed by the present invention.

20 Truncated elk-L, including soluble polypeptides, may be prepared by any of a number of conventional techniques. A desired DNA sequence may be chemically synthesized using known techniques. DNA fragments also may be produced by restriction endonuclease digestion of a full length cloned DNA sequence, and isolated by electrophoresis on agarose gels. Linkers containing restriction endonuclease
25 cleavage site(s) may be employed to insert the desired DNA fragment into an expression vector, or the fragment may be digested at cleavage sites naturally present therein. The well known polymerase chain reaction procedure also may be employed to isolate a DNA sequence encoding a desired protein fragment. As a further alternative, known mutagenesis techniques may be employed to insert a stop codon at a desired
30 point, e.g., immediately downstream of the codon for the last amino acid of the extracellular domain.

 In another approach, enzymatic treatment (e.g., using Bal 31 exonuclease) may be employed to delete terminal nucleotides from a DNA fragment to obtain a fragment having a particular desired terminus. Among the commercially available linkers are
35 those that can be ligated to the blunt ends produced by Bal 31 digestion, and which contain restriction endonuclease cleavage site(s). Alternatively, oligonucleotides that reconstruct the N- or C-terminus of a DNA fragment to a desired point may be

synthesized. The oligonucleotide may contain a restriction endonuclease cleavage site upstream of the desired coding sequence and position an initiation codon (ATG) at the N-terminus of the coding sequence.

5 The present invention provides purified elk-L polypeptides, both recombinant and non-recombinant. Variants and derivatives of native elk-L proteins that retain the desired biological activity (e.g., the ability to bind elk) are also within the scope of the present invention. elk-L variants may be obtained by mutations of nucleotide sequences coding for native elk-L polypeptides. An elk-L variant, as referred to herein, is a polypeptide substantially homologous to a native elk-L, but which has an amino
10 acid sequence different from that of native elk-L (human, murine or other mammalian species) because of one or more deletions, insertions or substitutions.

The variant amino acid sequence preferably is at least 80% identical to a native elk-L amino acid sequence, most preferably at least 90% identical. The percent identity may be determined, for example, by comparing sequence information using the GAP
15 computer program, version 6.0 described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). The preferred default parameters for the GAP program include:
20 (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional
25 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps. For elk-L fragments, the percent identity is based on that portion of a native elk-L protein that is present in the fragment.

Alterations of the native amino acid sequence may be accomplished by any of a number of known techniques. Mutations can be introduced at particular loci by
30 synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can
35 be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et

al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); Kunkel (*Proc. Natl. Acad. Sci. USA* 82:488, 1985); Kunkel et al. (*Methods in Enzymol.* 154:367, 1987); and U.S. Patent Nos. 4,518,584 and 4,737,462, which are incorporated by
5 reference herein.

Variants may comprise conservatively substituted sequences, meaning that a given amino acid residue is replaced by a residue having similar physiochemical characteristics. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or
10 substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known.

elk-L also may be modified to create elk-L derivatives by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives of elk-L may be prepared
15 by linking the chemical moieties to functional groups on elk-L amino acid side chains or at the N-terminus or C-terminus of a elk-L polypeptide or the extracellular domain thereof. Other derivatives of elk-L within the scope of this invention include covalent or aggregative conjugates of elk-L or its fragments with other proteins or polypeptides,
20 such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugate may comprise a signal or leader polypeptide sequence (e.g. the α -factor leader of *Saccharomyces*) at the N-terminus of a elk-L polypeptide. The signal or leader peptide co-translationally or post-translationally directs transfer of the conjugate from its side of synthesis to a site inside or outside of the cell membrane or
25 cell wall.

elk-L polypeptide fusions can comprise peptides added to facilitate purification and identification of elk-L. Such peptides include, for example, poly-His or the antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *Bio/Technology* 6:1204, 1988. One such peptide is the FLAG[®] peptide, Asp-
30 Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK), which is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody enabling rapid assay and facile purification of expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing. Fusion proteins capped with this peptide may also be
35 resistant to intracellular degradation in *E. coli*. A murine hybridoma designated 4E11 produces a monoclonal antibody that binds the peptide DYKDDDDK in the presence of certain divalent metal cations (as described in U.S. Patent 5,011,912, hereby

incorporated by reference) and has been deposited with the American Type Culture Collection under accession no HB 9259.

The present invention further includes elk-L polypeptides with or without associated native-pattern glycosylation. elk-L expressed in yeast or mammalian expression systems (e.g., COS-7 cells) may be similar to or significantly different from a native elk-L polypeptide in molecular weight and glycosylation pattern, depending upon the choice of expression system. Expression of elk-L polypeptides in bacterial expression systems, such as *E. coli*, provides non-glycosylated molecules.

DNA constructs that encode various additions or substitutions of amino acid residues or sequences, or deletions of terminal or internal residues or sequences not needed for biological activity or binding can be prepared. For example, N-glycosylation sites in the elk-L extracellular domain can be modified to preclude glycosylation, allowing expression of a more homogeneous, reduced carbohydrate analog in mammalian and yeast expression systems. N-glycosylation sites in eukaryotic polypeptides are characterized by an amino acid triplet Asn-X-Y, wherein X is any amino acid except Pro and Y is Ser or Thr. The human elk-L protein comprises one such triplet, at amino acids 115-117 of SEQ ID NO:2. Appropriate modifications to the nucleotide sequence encoding this triplet will result in substitutions, additions or deletions that prevent attachment of carbohydrate residues at the Asn side chain. Alteration of a single nucleotide, chosen so that Asn is replaced by a different amino acid, for example, is sufficient to inactivate an N-glycosylation site. Known procedures for inactivating N-glycosylation sites in proteins include those described in U.S. Patent 5,071,972 and EP 276,846, hereby incorporated by reference.

In another example, sequences encoding Cys residues that are not essential for biological activity can be altered to cause the Cys residues to be deleted or replaced with other amino acids, preventing formation of incorrect intramolecular disulfide bridges upon renaturation. Other variants are prepared by modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. EP 212,914 discloses the use of site-specific mutagenesis to inactivate KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Lys-Lys pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites. Human elk-L contains three KEX2 protease processing sites at amino acids 242-243, 243-244, and 246-247 of SEQ ID NO:2.

Naturally occurring elk-L variants are also encompassed by the present invention. Examples of such variants are proteins that result from alternative mRNA splicing events or from proteolytic cleavage of the elk-L protein, wherein the elk-binding property is retained. Alternative splicing of mRNA may yield a truncated but
5 biologically active elk-L protein, such as a naturally occurring soluble form of the protein, for example. Variations attributable to proteolysis include, for example, differences in the N- or C-termini upon expression in different types of host cells, due to proteolytic removal of one or more terminal amino acids from the elk-L protein (generally from 1-5 terminal amino acids).

10 Nucleic acid sequences within the scope of the present invention include isolated DNA and RNA sequences that hybridize to the native elk-L nucleotide sequences disclosed herein under conditions of moderate or severe stringency, and which encode biologically active elk-L. Moderate stringency hybridization conditions refer to conditions described in, for example, Sambrook et al. *Molecular Cloning: A*
15 *Laboratory Manual*, 2 ed. Vol. 1, pp. 1.101-104, Cold Spring Harbor Laboratory Press, (1989). Conditions of moderate stringency, as defined by Sambrook et al., include use of a prewashing solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0) and hybridization conditions of about 55°C, 5 X SSC, overnight. Conditions of severe stringency include higher temperatures of hybridization and washing. The skilled
20 artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as the length of the probe.

Due to the known degeneracy of the genetic code wherein more than one codon can encode the same amino acid, a DNA sequence may vary from that presented in SEQ ID NO:1, and still encode an elk-L protein having the amino acid sequence of SEQ ID
25 NO:1. Such variant DNA sequences may result from silent mutations (e.g., occurring during PCR amplification), and may be the product of deliberate mutagenesis of a native sequence.

The present invention thus provides isolated DNA sequences encoding biologically active elk-L, selected from: (a) DNA derived from the coding region of a
30 native mammalian elk-L gene (e.g., cDNA comprising the nucleotide sequence presented in SEQ ID NO:1); (b) DNA capable of hybridization to a DNA of (a) under moderately stringent conditions and which encodes biologically active elk-L; and (c) DNA which is degenerate as a result of the genetic code to a DNA defined in (a) or (b) and which encodes biologically active elk-L. The elk-L proteins encoded by such DNA
35 sequences are encompassed by the present invention.

Examples of elk-L proteins encoded by DNA that varies from the native DNA sequence of SEQ ID NO:1, wherein the variant DNA will hybridize to the native DNA

sequence under moderately stringent conditions, include, but are not limited to, elk-L fragments (soluble or membrane-bound) and elk-L proteins comprising inactivated N-glycosylation site(s), inactivated KEX2 protease processing site(s), or conservative amino acid substitution(s), as described above. Elk-L proteins encoded by DNA driven
5 from other mammalian species, wherein the DNA will hybridize to the human DNA of SEQ ID NO:1, are also encompassed.

Variants possessing the requisite ability to bind elk may be identified by any suitable assay. Biological activity of elk-L may be determined, for example, by competition for binding to the ligand binding domain of elk (i.e. competitive binding
10 assays).

One type of a competitive binding assay for elk-L polypeptide uses a radiolabeled, soluble human elk-L and intact cells expressing cell surface elk. Instead of intact cells, one could substitute soluble elk (such as an elk/Fc fusion protein) bound to a solid phase through a Protein A or Protein G interaction with the Fc region of the
15 fusion protein. Another type of competitive binding assay utilizes radiolabeled soluble elk such as an elk/Fc fusion protein, and intact cells expressing elk-L. Alternatively, soluble elk-L could be bound to a solid phase.

Competitive binding assays can be performed using standard methodology. For example, radiolabeled elk-L can be used to compete with a putative elk-L homolog
20 to assay for binding activity against surface-bound elk. Qualitative results can be obtained by competitive autoradiographic plate binding assays, or Scatchard plots may be utilized to generate quantitative results.

Alternatively, soluble elk can be bound to a solid phase such as a column chromatography matrix or a similar substrate suitable for analysis for the presence of a
25 detectable moiety such as ^{125}I . Binding to a solid phase can be accomplished, for example, by binding an elk/Fc fusion protein to a protein A or protein G-containing matrix.

The binding characteristics of elk-L (including variants) may also be determined using the conjugated, soluble elk (for example, ^{125}I -elk/Fc) in competition assays
30 similar to those described above. In this case, however, intact cells expressing elk-L, or soluble elk-L bound to a solid substrate, are used to measure the extent to which a sample containing a putative elk variant competes for binding of a conjugated soluble elk to elk-L.

The elk-L of the present invention can be used in a binding assay to detect cells
35 expressing elk. For example, elk-L or an extracellular domain or a fragment thereof can be conjugated to a detectable moiety such as ^{125}I . Radiolabeling with ^{125}I can be performed by any of several standard methodologies that yield a functional ^{125}I -elk-L

molecule labeled to high specific activity. Alternatively, another detectable moiety such as an enzyme that can catalyze a colorimetric or fluorometric reaction, biotin or avidin may be used. Cells to be tested for elk expression can be contacted with labeled elk-L. After incubation, unbound labeled elk-L is removed and binding is measured using the
5 detectable moiety.

The elk ligand proteins disclosed herein also may be employed to measure the biological activity of elk protein in terms of binding affinity for elk-L. To illustrate, elk-L may be employed in a binding affinity study to measure the biological activity of an elk protein that has been stored at different temperatures, or produced in different
10 cell types. The biological activity of an elk protein thus can be ascertained before it is used in a research study, for example.

Elk-L proteins find use as reagents that may be employed by those conducting "quality assurance" studies, e.g., to monitor shelf life and stability of elk protein under different conditions. Elk ligands may be used in determining whether biological
15 activity is retained after modification of a elk protein (e.g., chemical modification, truncation, mutation, etc.). The binding affinity of the modified elk protein for an elk-L is compared to that of an unmodified elk protein to detect any adverse impact of the modifications on biological activity of elk.

A different use of an elk ligand is as a reagent in protein purification
20 procedures. Elk-L or elk-L/Fc fusion proteins may be attached to a solid support material by conventional techniques and used to purify elk by affinity chromatography.

elk-L polypeptides may exist as oligomers, such as dimers or trimers. Oligomers are linked by disulfide bonds formed between cysteine residues on different elk-L polypeptides. In one embodiment of the invention, an elk-L dimer is created by
25 fusing elk-L to the Fc region of an antibody (IgG1) in a manner that does not interfere with binding of elk-L to the elk ligand binding domain. The Fc polypeptide preferably is fused to the C-terminus of a soluble elk-L (comprising only the extracellular domain). Preparation of fusion proteins comprising heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been
30 described, e.g., by Ashkenazi et al. (*PNAS USA* 88:10535, 1991) and Byrn et al. (*Nature* 344:677, 1990), hereby incorporated by reference. A gene fusion encoding the elk-L/Fc fusion protein is inserted into an appropriate expression vector. The elk-L/Fc fusion proteins are allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between Fc polypeptides, yielding divalent elk-L. If
35 fusion proteins are made with both heavy and light chains of an antibody, it is possible to form a elk-L oligomer with as many as four elk-L extracellular regions.

Alternatively, one can link two soluble elk-L domains with a peptide linker such as the Gly4SerGly5Ser linker sequence described in United States Patent 5,073,627.

The present invention provides oligomers of elk-L extracellular domains or fragments thereof, linked by disulfide interactions, or expressed as fusion polymers with or without spacer amino acid linking groups. For example, a dimer of the elk-L extracellular domain can be linked by an IgG Fc region linking group.

The present invention provides recombinant expression vectors for expression of elk-L, and host cells transformed with the expression vectors. Any suitable expression system may be employed. The vectors include an elk-L DNA sequence operably linked to suitable transcriptional or translational regulatory nucleotide sequences, such as those derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, an mRNA ribosomal binding site, and appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the elk-L DNA sequence. Thus, a promoter nucleotide sequence is operably linked to a elk-L DNA sequence if the promoter nucleotide sequence controls the transcription of the elk-L DNA sequence. The ability to replicate in the desired host cells, usually conferred by an origin of replication, and a selection gene by which transformants are identified, may additionally be incorporated into the expression vector.

In addition, sequences encoding appropriate signal peptides that are not native to the elk-L gene can be incorporated into expression vectors. For example, a DNA sequence for a signal peptide (secretory leader) may be fused in frame to the elk-L sequence so that the elk-L is initially translated as a fusion protein comprising the signal peptide. A signal peptide that is functional in the intended host cells enhances extracellular secretion of the elk-L polypeptide. The signal peptide is cleaved from the elk-L polypeptide upon secretion of elk-L from the cell.

Suitable host cells for expression of elk-L polypeptides include prokaryotes, yeast or higher eukaryotic cells. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985). Cell-free translation systems could also be employed to produce elk-L polypeptides using RNAs derived from DNA constructs disclosed herein.

Prokaryotes include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a prokaryotic host cell,

such as *E. coli*, an elk-L polypeptide may include an N-terminal methionine residue to facilitate expression of the recombinant polypeptide in the prokaryotic host cell. The N-terminal Met may be cleaved from the expressed recombinant elk-L polypeptide.

Expression vectors for use in prokaryotic host cells generally comprise one or
5 more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline
10 resistance and thus provides simple means for identifying transformed cells. An appropriate promoter and a elk-L DNA sequence are inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA).

Promoter sequences commonly used for recombinant prokaryotic host cell
15 expression vectors include β -lactamase (penicillinase), lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EP-A-36776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful prokaryotic host cell
20 expression system employs a phage λ P_L promoter and a cI857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2 (resident in *E. coli* strain JMB9 (ATCC 37092)) and pPLc28 (resident in *E. coli* RR1 (ATCC 53082)).

25 elk-L alternatively may be expressed in yeast host cells, preferably from the *Saccharomyces* genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia* or *Kluyveromyces*, may also be employed. Yeast vectors will often contain an origin of replication sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription
30 termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-
35 phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other

suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA-73,657. Another alternative is the glucose-repressible ADH2 promoter described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). Shuttle vectors replicable in both yeast and *E. coli* may be

- 5 constructed by inserting DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) into the above-described yeast vectors.

The yeast α -factor leader sequence may be employed to direct secretion of the elk-L polypeptide. The α -factor leader sequence is often inserted between the promoter sequence and the structural gene sequence. See, e.g., Kurjan et al., *Cell* 30:933, 1982; 10 Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984; U. S. Patent 4,546,082; and EP 324,274. Other leader sequences suitable for facilitating secretion of recombinant polypeptides from yeast hosts are known to those of skill in the art. A leader sequence may be modified near its 3' end to contain one or more restriction sites. This will facilitate fusion of the leader sequence to the structural gene.

- 15 Yeast transformation protocols are known to those of skill in the art. One such protocol is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978. The Hinnen et al. protocol selects for Trp⁺ transformants in a selective medium, wherein the selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 μ g/ml adenine and 20 μ g/ml uracil.

- 20 Yeast host cells transformed by vectors containing ADH2 promoter sequence may be grown for inducing expression in a "rich" medium. An example of a rich medium is one consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 μ g/ml adenine and 80 μ g/ml uracil. Derepression of the ADH2 promoter occurs when glucose is exhausted from the medium.

- 25 Mammalian or insect host cell culture systems could also be employed to express recombinant elk-L polypeptides. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *BioTechnology* 6:47 (1988). Established cell lines of mammalian origin also may be employed. Examples of suitable mammalian host cell lines include the COS-7 line of 30 monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, and BHK (ATCC CRL 10) cell lines, and the CV-1/EBNA-1 cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10: 2821, 1991).

- 35 Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2,

Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* I site located in the SV40 viral origin of replication site is included.

Exemplary expression vectors for use in mammalian host cells can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983). A useful system for stable high level expression of mammalian cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A useful high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984 has been deposited as ATCC 39890. Additional useful mammalian expression vectors are described in EP-A-0367566, and in U.S. Patent Application Serial No. 07/701,415, filed May 16, 1991, incorporated by reference herein. The vectors may be derived from retroviruses. In place of the native signal sequence, a heterologous signal sequence may be added, such as the signal sequence for interleukin-7 (IL-7) described in United States Patent 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., *Nature* 312:768 (1984); the interleukin-4 signal peptide described in EP 367,566; the type I interleukin-1 receptor signal peptide described in U.S. Patent 4,968,607; and the type II interleukin-1 receptor signal peptide described in EP 460,846.

The present invention provides substantially homogeneous elk-L protein, which may be produced by recombinant expression systems as described above or purified from naturally occurring cells. The elk-L is purified to substantial homogeneity, as indicated by a single protein band upon analysis by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

One process for producing the elk-L protein comprises culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes elk-L under conditions such that elk-L is expressed. The elk-L protein is then recovered from culture medium or cell extracts, depending upon the expression system employed. As the skilled artisan will recognize, procedures for purifying the recombinant elk-L will vary according to such factors as the type of host cells employed and whether or not the elk-L is secreted into the culture medium.

For example, when expression systems that secrete the recombinant protein are employed, the culture medium first may be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred. Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, (e.g., silica gel having pendant methyl or other aliphatic groups) can be employed to further purify elk-L. Some or all of the foregoing purification steps, in various combinations, can be employed to provide a substantially homogeneous recombinant protein.

It is also possible to utilize an affinity column comprising the ligand binding domain of elk to affinity-purify expressed elk-L polypeptides. elk-L polypeptides can be removed from an affinity column in a high salt elution buffer and then dialyzed into a lower salt buffer for use. Alternatively, the affinity column may comprise an antibody that binds elk-L. Example 4 describes a procedure for employing the elk-L protein of the present invention to generate monoclonal antibodies directed against elk-L.

Recombinant protein produced in bacterial culture is usually isolated by initial disruption of the host cells, centrifugation, extraction from cell pellets if an insoluble polypeptide, or from the supernatant fluid if a soluble polypeptide, followed by one or more concentration, salting-out, ion exchange, affinity purification or size exclusion chromatography steps. Finally, RP-HPLC can be employed for final purification steps. Microbial cells can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Transformed yeast host cells are preferably employed to express elk-L as a secreted polypeptide. This simplifies purification. Secreted recombinant polypeptide from a yeast host cell fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). Urdal et al. describe two sequential, reversed-phase HPLC steps for purification of recombinant human IL-2 on a preparative HPLC column.

Nucleic Acid Fragments

The present invention further provides fragments of the elk-L nucleotide sequences presented herein. Such fragments desirably comprise at least about 14 nucleotides of the sequence presented in SEQ ID NO:1. DNA and RNA complements
5 of said fragments are provided herein, along with both single-stranded and double-stranded forms of the elk-L DNA.

Among the uses of such elk-L nucleic acid fragments is use as a probe. Such probes may be employed in cross-species hybridization procedures to isolate elk-L DNA from additional mammalian species. As one example, a probe
10 corresponding to the extracellular domain of elk-L may be employed. The probes also find use in detecting the presence of elk-L nucleic acids in *in vitro* assays and in such procedures as Northern and Southern blots. Cell types expressing elk-L can be identified. Such procedures are well known, and the skilled artisan can choose a probe of suitable length, depending on the particular intended application.

15 Other useful fragments of the elk-L nucleic acids are antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target elk-L mRNA (sense) or elk-L DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of elk-L cDNA. Such a fragment generally
20 comprises at least about 14 nucleotides, preferably from about 14 to about 30 nucleotides. The ability to create an antisense or a sense oligonucleotide, based upon a cDNA sequence for a given protein is described in, for example, Stein and Cohen, *Cancer Res.* 48:2659, 1988 and van der Krol et al., *BioTechniques* 6:958, 1988.

Binding of antisense or sense oligonucleotides to target nucleic acid sequences
25 results in the formation of duplexes that block translation (RNA) or transcription (DNA) by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of elk-L proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-
30 phosphodiester backbones (or other sugar linkages, such as those described in WO91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences. Other examples of sense or antisense oligonucleotides
35 include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10448, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further

still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides may be introduced into a cell
5 containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO₄-mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. Antisense or sense oligonucleotides are preferably introduced into a cell containing the target nucleic acid sequence by insertion of the antisense or sense oligonucleotide into a suitable retroviral vector, then contacting the
10 cell with the retrovirus vector containing the inserted sequence, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see PCT Application US 90/02656).

Sense or antisense oligonucleotides also may be introduced into a cell
15 containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand
20 binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid
25 complex is preferably dissociated within the cell by an endogenous lipase.

The following examples are provided to illustrate particular embodiments and not to limit the scope of the invention.

EXAMPLE 1

Preparation of Soluble elk/Fc Fusion Protein

This example describes construction of an expression vector encoding a soluble elk/Fc fusion protein, for use in detecting cDNA clones encoding an elk ligand (elk-L). A DNA and encoded amino acid sequence for rat elk cDNA is presented in Lhotak et al. (*Mol. Cell. Biol.* 11:2496, 1991), hereby incorporated by reference. The rat elk
35 protein has a 538 amino acid extracellular domain, a 25 amino acid transmembrane domain, and a 419 amino acid cytoplasmic domain.

A rat'elk cDNA fragment was fused to the N-terminus of cDNA encoding the Fc portion of a human IgG1 antibody. The rat elk cDNA was obtained from T. Pawson (Samuel Lunenfeld Research Institute, Mt. Sinai Hospital, Toronto). An Asp718 restriction endonuclease cleavage site was introduced upstream of the elk coding region. An Asp 718-BglII fragment of rat elk cDNA (comprising the entire extracellular domain, the transmembrane region, and a small portion of the cytoplasmic domain) was isolated.

DNA encoding a single chain polypeptide comprising the Fc region of a human IgG1 antibody was cloned into the *SpeI* site of the pBLUESCRIPT SK[®] vector, which is commercially available from Stratagene Cloning Systems, La Jolla, California. This plasmid vector is replicable in *E. coli* and contains a polylinker segment that includes 21 unique restriction sites. A unique *BglII* site was introduced near the 5' end of the inserted Fc encoding sequence, such that the *BglII* site encompasses the codons for amino acids three and four of the Fc polypeptide.

The encoded Fc polypeptide extends from the N-terminal hinge region to the native C-terminus, i.e., is an essentially full-length antibody Fc region. Fragments of Fc regions, e.g., those that are truncated at the C-terminal end, also may be employed. The fragments preferably contain multiple cysteine residues (at least the cysteine residues in the hinge reaction) to permit interchain disulfide bonds to form between the Fc polypeptide portions of two separate elk/Fc fusion proteins, forming dimers as discussed above.

The above-described Asp718-BglII elk cDNA fragment was cloned into the pBLUESCRIPT SK[®] vector containing the Fc cDNA, such that the elk cDNA is positioned upstream of the Fc cDNA. Single stranded DNA derived from the resulting gene fusion was mutagenized by the method described in Kunkel (*Proc. Natl. Acad. Sci. USA* 82:488, 1985) and Kunkel et al. (*Methods in Enzymol.* 154:367, 1987) in order to perfectly fuse the entire extracellular domain of elk to the Fc sequence. The mutagenized DNA was sequenced to confirm that the proper nucleotides had been removed (i.e., transmembrane region and partial cytoplasmic domain DNA was deleted) and that the elk and Fc sequences were in the same reading frame. The fusion cDNA was then excised and inserted into a mammalian expression vector designated pCAV/DHFR. The pCAV/DHFR vector is similar to pCAV/NOT, described in PCT Application WO 90/05183, but additionally contains a dihydrofolate reductase (DHFR) gene as a selectable marker. The DHFR gene confers a selective advantage on otherwise DHFR⁻ mammalian cells that have taken up the vector, when grown in the presence of methotrexate (MTX).

elk/Fc fusion proteins preferably are synthesized in recombinant mammalian cell culture. The elk/Fc fusion-containing expression vector was transfected into CV-1 cells (ATCC CCL 70) and COS-7 cells (ATCC CRL 1651), both derived from monkey kidney. The elk/Fc fusion protein expression level was relatively low in both CV-1 and
5 COS-7 cells. Thus, expression in 293 cells (transformed primary human embryonal kidney cells, ATCC CRL 1573) was attempted.

The elk-Fc gene fusion was inserted into a mammalian expression vector designated HAV-EO (Dower et al., *J. Immunol.* 142:4314, 1989). 293 cells transfected with the HAV-EO/elk/Fc vector were cultivated in roller bottles to allow
10 transient expression of the fusion protein, which is secreted into the culture medium *via* the elk signal peptide. The fusion protein was purified on protein A sepharose columns, eluted, and used to screen cells for the ability to bind the elk/Fc protein, as described in examples 2 and 3.

15 **EXAMPLE 2: Screening Cells for elk/Fc Binding**

Various cell types were screened for the ability to bind elk/Fc, to identify candidate cell types useful as nucleic acid sources in an attempt to clone an elk ligand. The recombinant rat elk/Fc protein produced and purified in Example 1 was
20 radiolabeled using a commercially available Enzymobead radioiodination reagent (BioRad) according to manufacturer's instructions. The ^{125}I -elk/Fc protein did not show specific binding to any of several cell lines tested. Concern that direct labeling of the elk/Fc protein with Na^{125}I may have resulted in loss of biological activity led to the development of a two-step binding assay.

The two-step assay involved incubating cells with non-radiolabeled elk/Fc
25 followed by ^{125}I -mouse anti-human Fc antibody. The latter antibody will bind to the Fc portion of any elk/Fc fusion protein that has bound to the cells.

The mouse anti-human Fc antibody was obtained from Jackson Immunoresearch Laboratories, Inc., West Grove, PA. This antibody showed minimal binding to Fc proteins bound to the Fc γ receptor. The antibody was labeled using the
30 Chloramine T method. Briefly, a P6 column was prepared according to the manufacturer's instructions. In a microfuge tube, 10 μg of antibody was dissolved in 10 μl of PBS. 2000 μCi of carrier-free Na^{125}I was added and the solution was mixed well. 15 μl of a freshly prepared solution of chloramine-T (32 $\mu\text{g}/\text{ml}$ in 0.05 M sodium phosphate buffer (pH 7.2) was then added and the mixture was incubated for 30
35 minutes at room temperature. The mixture was immediately applied to the P6 column. The radiolabelled antibody was then eluted from the column by collecting 100-150 μl fractions of eluate. Binding media (RPMI 1640 containing 25 mg/ml bovine serum

albumin, 2 mg/ml sodium azide, 20 mM Hepes pH 7.2) was added to peak fractions to bring the total volume of each fraction to 2 ml. Radioiodination yielded specific activities in the range of $5-10 \times 10^{15}$ cpm/mmol protein.

The screening procedure was as follows. Approximately 2×10^6 cells were
5 cultured in 96-well plates. 150 μ l binding medium was added to each well and the cells were then incubated in the presence or absence of elk/Fc for 1 hour at 37°C with gentle agitation. The cells were sedimented from the mixture by centrifugation of the 96-well plates, washed with PBS, centrifuged again, and resuspended in binding medium. The cells were then incubated with the 125 I-mouse anti-human Fc antibody,
10 prepared as described above, for 1 hour at 37°C. Cells were also incubated with 125 I-mouse anti-human Fc antibody in the presence of excess unlabelled anti-human Fc antibody as a negative control. Following a one hour incubation with the 125 I-antibody, cells and unbound 125 I-antibody were separated by the phthalate oil separation method, essentially as described by Dower et al., *J. Immunol.* 132:751,
15 1984. Cell-bound and free 125 I-antibody were quantitated on a Packard Autogamma counter. No cell types were found to bind elk/Fc in this assay.

Cells, including additional cell types not previously screened, were assayed for elk/Fc binding by a third procedure. Cells were incubated with elk/Fc, followed by a biotinylated anti-human Fc antibody, followed by streptavidin-phycoerythrin (Becton
20 Dickinson). The biotinylated antibody was purchased from Jackson Immunoresearch Laboratories. Streptavidin binds to the biotin molecule attached to the anti-human Fc antibody, which in turn binds to the Fc portion of the elk/Fc fusion protein. Phycoerythrin is a fluorescent phycobiliprotein which serves as a detectable label. The level of fluorescence signal was measured for each cell type using a FACScan® flow
25 cytometer (Becton Dickinson).

The cells screened for the ability to bind elk/Fc by at least one of the three procedures described above included approximately 30 different cell types falling into the following general categories: primary murine fetal brain cells, murine fetal liver cell lines, murine fetal brain cell lines, and human lung carcinoma (fibroblastoid) cell lines.
30 No cell types deemed positive for elk/Fc binding were identified.

EXAMPLE 3: Isolation of Elk Ligand cDNA from Human Placental cDNA Library

None of the cell types assayed in Example 2 exhibited detectable elk/Fc binding.
35 A human placental cDNA library was chosen as a possible source of elk ligand cDNA, since placental tissue is rich in growth and differentiation factors. As discussed above, it is likely that the elk ligand plays a role in growth, survival, or differentiation of cells

expressing elk (the corresponding receptor). However, the fact that placenta is relatively rich in Fc receptors posed a potential problem for the cloning attempt, since an elk/Fc fusion was used to screen clones.

A human placental cDNA library in plasmid pDC302 was prepared as described
5 in Larsen et al. (*J. Exp. Med.*, 172:1559, 1990). pDC302 is a mammalian expression vector that also replicates in *E. coli* (Mosley et al., *Cell* 59:335, 1989). Briefly, placental cDNA was cloned into the *Bgl*III site of pDC302 by an adaptor method similar to that described by Haymerle et al (*Nucl. Acids Res.* 14:8615, 1986).

E. coli strain DH5 α cells transfected with the placental cDNA library in pDC302
10 were plated to provide approximately 500 colonies per plate. Colonies were scraped from each plate, pooled, and plasmid DNA prepared from each pool. The pooled DNA representing about 2000 colonies was then used to transfect a sub-confluent layer of CV-1/EBNA-1 cells using DEAE-dextran followed by chloroquine treatment, similar to that described by Luthman et al., *Nucl. Acids Res.* 11:1295, 1983) and McCutchan et al., *J. Natl. Cancer Inst.* 41:351, 1986). The CV-1/EBNA-1 cell line (ATCC
15 CRL10478) constitutively expresses EBV nuclear antigen-1 driven from the CMV immediate-early enhancer/promoter. CV1-EBNA-1 was derived from the African Green Monkey kidney cell line CV-1 (ATCC CCL 70), as described by McMahan et al. (*EMBO J.* 10:2821, 1991).

20 In order to transfect the CV-1/EBNA-1 cells with the cDNA library, the cells were maintained in complete medium (Dulbecco's modified Eagle's media (DMEM) containing 10% (v/v) fetal calf serum (FCS), 50 U/ml penicillin, 50 U/ml streptomycin, 2 mM L-glutamine) and were plated at a density of 2×10^5 cells/well on single-well chambered slides (Lab-Tek). Slides were pretreated with 1 ml human fibronectin (10
25 μ g/ml in PBS) for 30 minutes followed by 1 wash with PBS. Media was removed from the adherent cell layer and replaced with 1.5 ml complete medium containing 66.6 μ M chloroquine sulfate. 0.2 mls of DNA solution (2 μ g DNA, 0.5 mg/ml DEAE-dextran in complete medium containing chloroquine) was then added to the cells and incubated for 5 hours. Following the incubation, the media was removed and the cells
30 shocked by addition of complete medium containing 10% DMSO for 2.5 to 20 minutes followed by replacement of the solution with fresh complete medium. The cells were cultured for 2 to 3 days to permit transient expression of the inserted sequences.

Transfected monolayers of CV-1/EBNA-1 cells were assayed for expression of elk-L by slide autoradiography essentially as described by Gearing et al. (*EMBO J.*
35 8:3667, 1989). Transfected CV-1/EBNA-1 cells (adhered to chambered slides) were washed once with binding medium with nonfat dry milk (BM-NFDM) (RPMI medium 1640 containing 25 mg/ml bovine serum albumin (BSA), 2 mg/ml sodium azide, 20

mM HEPES, pH 7.2, and 50 mg/ml nonfat dry milk). Cells were then incubated with elk/Fc in BM-NFDM (1 µg/ml) for 1 hour at room temperature. After incubation, the cell monolayers in the chambered slides were washed three times with BM-NFDM to remove unbound elk/Fc fusion protein and then incubated with 40 ng/ml ¹²⁵I-mouse anti-human Fc antibody prepared in Example 2 (a 1:50 dilution) for 1 hour at room temperature. The cells were washed three times with BM-NFDM, followed by 2 washes with phosphate-buffered saline (PBS) to remove unbound ¹²⁵I-mouse anti-human Fc antibody. The cells were fixed by incubating for 30 minutes at room temperature in 2.5% glutaraldehyde in PBS, pH 7.3, washed twice in PBS and air dried. The chamber slides containing the cells were exposed on a Phosphorimager (Molecular Dynamics) overnight, then dipped in Kodak GTNB-2 photographic emulsion (6x dilution in water) and exposed in the dark for 3-5 days at 4°C in a light proof box. The slides were then developed for approximately 4 minutes in Kodak D19 developer (40 g/500 ml water), rinsed in water and fixed in Agfa G433C fixer. The slides were individually examined with a microscope at 25-40x magnification and positive cells expressing elk-L were identified by the presence of autoradiographic silver grains against a light background.

Using the slide autoradiography approach, approximately 300,000 cDNAs were screened in pools of approximately 2,000 cDNAs until assay of one transfectant pool showed multiple cells clearly positive for elk/Fc binding. This pool was then partitioned into pools of 500 and again screened by slide autoradiography and a positive pool was identified. Individual colonies from this pool of 500 were screened until a single clone (clone #E7) was identified which directed synthesis of a surface protein with detectable elk/Fc binding activity. This clone was isolated, and its cDNA insert was sequenced.

The nucleotide and encoded amino acid sequences of the coding region of the human elk ligand cDNA of clone E7 are presented in SEQ ID NOS:1 and 2. The cDNA insert is 2111 bp in length. There are three reasonably sized open reading frames within this sequence that could encode proteins of 82, 85, and 346 amino acids. Of these three, the two smallest open reading frames lack both good translation initiation codons as well as a signal sequence. Thus, DNA and encoded amino acid sequences for the 346-amino acid open reading frame are presented in SEQ ID NOS:1 and 2. The protein of SEQ ID NO:2 is a type I transmembrane protein, with an N-terminal signal peptide (amino acids -24 to -1, an extracellular domain (amino acids 1-213) a transmembrane domain (amino acids 214-241) and a cytoplasmic domain (amino acids 242-322). The predicted molecular weight of the native protein following cleavage of the signal sequence is 35,180 daltons. The mature protein has an estimated pI of

9.006. There are 307 bp of 5' noncoding sequence and 763 bp of 3' non-coding sequence flanking the coding region.

Human elk-L cDNA was excised from clone E7 by digestion with *Bsu*361 (which cleaves downstream of the stop codon), followed by treatment with Klenow
5 fragment to render the *Bsu*361-digested end blunt, followed by digestion with *Sma*I (which cleaves upstream of the initiation codon and generates blunt ends). The excised cDNA, which includes the entire coding region, was cloned into the *Sma*I site (in the multiple cloning site) of pBLUESCRIPT® SK(-) (Stratagene Cloning Systems, La Jolla, CA). The resulting vector (designated tele 7 in pBLUESCRIPT SK(-)) in *E. coli*
10 DH5α cells was deposited with the American Type Culture Collection, Rockville, MD, USA (ATCC) on October 9, 1992 and assigned accession number ATCC 69085. The deposit was made under the terms of the Budapest Treaty.

EXAMPLE 4: Monoclonal Antibodies to elk-L

15 This example illustrates the preparation of monoclonal antibodies to elk-L. elk-L is expressed in mammalian host cells such as COS-7 or CV-1/EBNA-1 cells and purified using elk/Fc affinity chromatography. Purified elk-L (or a fragment thereof such as the extracellular domain) can be used to generate monoclonal antibodies against elk-L using conventional techniques, for example, those techniques described in U.S.
20 Patent 4,411,993. Briefly, mice are immunized with elk-L as an immunogen emulsified in complete Freund's adjuvant, and injected in amounts ranging from 10-100 µg subcutaneously or intraperitoneally. Ten to twelve days later, the immunized animals are boosted with additional elk-L emulsified in incomplete Freund's adjuvant. Mice are periodically boosted thereafter on a weekly to bi-weekly immunization
25 schedule. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot blot assay or ELISA (Enzyme-Linked Immunosorbent Assay), for elk-L antibodies.

Following detection of an appropriate antibody titer, positive animals are provided one last intravenous injection of elk-L in saline. Three to four days later, the
30 animals are sacrificed, spleen cells harvested, and spleen cells are fused to a murine myeloma cell line, e.g., NS1 or preferably P3x63Ag8.653 (ATCC CRL 1580). Fusions generate hybridoma cells, which are plated in multiple microtiter plates in a HAT (hypoxanthine, aminopterin and thymidine) selective medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

35 The hybridoma cells are screened by ELISA for reactivity against purified elk-L by adaptations of the techniques disclosed in Engvall et al., *Immunochem.* 8:871, 1971 and in U.S. Patent 4,703,004. A preferred screening technique is the antibody capture

technique described in Beckmann et al., (*J. Immunol.* 144:4212, 1990) Positive hybridoma cells can be injected intraperitoneally into syngeneic BALB/c mice to produce ascites containing high concentrations of anti-elk-L monoclonal antibodies. Alternatively, hybridoma cells can be grown *in vitro* in flasks or roller bottles by various techniques. Monoclonal antibodies produced in mouse ascites can be purified by ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can also be used, as can affinity chromatography based upon binding to elk-L.

EXAMPLE 5: Northern Blot Analyses

To investigate expression of elk-L in various tissues, a human tissue Northern blot that contained 2 ug of poly A+ mRNA from a variety of tissues (available from Clontech, Palo Alto, CA) was probed with an elk-L riboprobe. 5' and 3' non-coding sequences were removed from the elk ligand cDNA by digesting with SmaI and BsuI and inserting the coding region into the pBluescript SK(-) vector (Stratagene Cloning Systems, LaJolla, CA). A riboprobe was prepared by cutting the elk-L cDNA with SfiI and using the T3 RNA polymerase promoter to transcribe the 3' most 450 nucleotides of elk-L coding sequence. The blot was hybridized with the riboprobe at 63° C, then washed, first with 1XSSC/0.1% SDS for 1 h at 68°C, then with 0.1X SSC/0.1%SDS for 30 min at 68°C. The blot was exposed to X-ray film at -70° for 11 days sandwiched between two intensifying screens. A major 3.5 kb message was observed in heart, placenta, lung, liver, skeletal muscle, kidney, and pancreas.

The elk-L mRNA was not present in detectable levels in whole adult brain, even though brain and testis are the only tissues reported to express the elk receptor (Lhotak et al., *supra*). It is possible that expression in brain is at very low levels, or is limited to a distinct region(s), and therefore was not detected by the foregoing technique. To further investigate this issue, dissected brains from 8-day old and young adult rats were probed with elk-L DNA. Elk-L appears to be expressed in the developing rat brain, with the highest level of expression in the olfactory bulb. Expression of elk-L in the olfactory bulb appears to continue into adulthood. Interestingly, the olfactory bulb is actively regenerating throughout the life of the animal, in contrast to the rest of the brain.

A second Northern blot revealed that elk-L mRNA expression is upregulated by TNF in human umbilical vein endothelial cells (HUVECS). B61 mRNA in HUVECS was shown to be inducible by treating the cells with IL-1 or TNF (Holzman et al., *supra*). In view of the degree of homology between elk-L and B61, a Northern blot of

HUVEC mRNA (from either resting cells or cells induced with TNF) was probed with the above-described elk-L riboprobe. The major 3.5 kb species seen in other tissues was specifically induced following treatment with TNF.

5 **EXAMPLE 6: Southern Blot Analysis**

The presence of elk-L DNA in various species was investigated as follows. A blot consisting of EcoRI-digested DNAs from a variety of species (Clontech, Palo Alto, CA) was prehybridized at 63°C in prehybridization buffer. The blot was then probed with a random primed clone E7 probe (corresponding to the entire elk-L coding region) and washed to high stringency (1XSSC/0.1% SDS for 60 min at 63°C, then 10 0.1XSSC/0.1% SDS at 63°C for 30 min). The image of the blot was developed on a Phosphorimager (Molecular Dynamics). The gene was found to be highly conserved in a wide variety of species, including monkey, rat, mouse, dog, cow, rabbit, chicken, and even yeast.

15 **EXAMPLE 7: Binding Study**

To examine the binding of elk-L to the elk receptor, a modified indirect Scatchard analysis was devised. This indirect assay was necessary because direct radiolabeling of elk/Fc inactivated the binding specificity of this protein. The assay procedure was as follows.

20 CV1-EBNA-1 cells (described in Example 3) in 12-well plates (2.5×10^5 cells/well) were transfected with an expression vector containing the clone E7 elk-L cDNA. The transfected cells were cultured for two days to permit expression of the elk-L protein, then washed with BM-NFDM (the binding medium described in Example 2, also containing 50 mg/ml nonfat dry milk), and incubated with various concentrations of the elk/Fc fusion protein prepared in Example 1, for 1 hour at 37°C. Subsequently, cells were washed and incubated with a constant saturating concentration of the ^{125}I -labeled mouse anti-human IgG antibody prepared in Example 2 (40 ng/ml) in binding medium with gentle agitation for 1 hour at 37°C. Cells then were harvested 25 by trypsinization. Cells and unbound ^{125}I -antibody were separated by a phthalate oil separation method, essentially as described by Dower et al., *J. Immunol.* 132:751, 1984.

30 Free and cell-bound ^{125}I -antibody were quantified on a Packard Autogamma Counter. The binding curve was replotted in the Scatchard coordinate system (Scatchard, *Ann. N.Y. Acad. Sci.* 51:660, 1949). Affinity calculations were generated on RS/1 (BBN Software, Boston, MA) run on a Microvax computer. A single high affinity binding component was identified. The elk-L binding sites expressed on the

CV-1 EBNA-1 transfectants had an affinity constant (K_A) of $1.08 \times 10^9 \text{ M}^{-1}$ for elk/Fc. This affinity constant matches well with those observed for binding of elk/Fc to the native ligand expressed on various rat neutral cell lines.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Lyman, Stewart D.
Beckmann, M. Patricia
Baum, Peter R
- (ii) TITLE OF INVENTION: Novel Cytokine Designated elk Ligand
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Immunex Corporation
 - (B) STREET: 51 University Street
 - (C) CITY: Seattle
 - (D) STATE: Washington
 - (E) COUNTRY: USA
 - (F) ZIP: 98101
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 12-NOV-1993
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Seese, Kathryn A.
 - (B) REGISTRATION NUMBER: 32,172
 - (C) REFERENCE/DOCKET NUMBER: 2807-WO
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (206) 587-0430
 - (B) TELEFAX: (206) 233-0644

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1041 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 1..1041

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 73..1038

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GCT CGG CCT GGG CAG CGT TGG CTC GGC AAG TGG CTT GTG GCG ATG	48
Met Ala Arg Pro Gly Gln Arg Trp Leu Gly Lys Trp Leu Val Ala Met	
-24 -20 -15 -10	
GTC GTG TGG GCG CTG TGC CGG CTC GCC ACA CCG CTG GCC AAG AAC CTG	96
Val Val Trp Ala Leu Cys Arg Leu Ala Thr Pro Leu Ala Lys Asn Leu	
-5 1 5	
GAG CCC GTA TCC TGG AGC TCC CTC AAC CCC AAG TTC CTG AGT GGG AAG	144
Glu Pro Val Ser Trp Ser Ser Leu Asn Pro Lys Phe Leu Ser Gly Lys	
10 15 20	
GGC TTG GTG ATC TAT CCG AAA ATT GGA GAC AAG CTG GAC ATC ATC TGC	192
Gly Leu Val Ile Tyr Pro Lys Ile Gly Asp Lys Leu Asp Ile Ile Cys	
25 30 35 40	
CCC CGA GCA GAA GCA GGG CGG CCC TAT GAG TAC TAC AAG CTG TAC CTG	240
Pro Arg Ala Glu Ala Gly Arg Pro Tyr Glu Tyr Tyr Lys Leu Tyr Leu	
45 50 55	
GTG CGG CCT GAG CAG GCA GCT GCC TGT AGC ACA GTT CTC GAC CCC AAC	288
Val Arg Pro Glu Gln Ala Ala Ala Cys Ser Thr Val Leu Asp Pro Asn	
60 65 70	
GTG TTG GTC ACC TGC AAT AGG CCA GAG CAG GAA ATA CGC TTT ACC ATC	336
Val Leu Val Thr Cys Asn Arg Pro Glu Gln Glu Ile Arg Phe Thr Ile	
75 80 85	
AAG TTC CAG GAG TTC AGC CCC AAC TAC ATG GGC CTG GAG TTC AAG AAG	384
Lys Phe Gln Glu Phe Ser Pro Asn Tyr Met Gly Leu Glu Phe Lys Lys	
90 95 100	
CAC CAT GAT TAC TAC ATT ACC TCA ACA TCC AAT GGA AGC CTG GAG GGG	432
His His Asp Tyr Tyr Ile Thr Ser Thr Ser Asn Gly Ser Leu Glu Gly	
105 110 115 120	
CTG GAA AAC CGG GAG GGC GGT GTG TGC CGC ACA CGC ACC ATG AAG ATC	480
Leu Glu Asn Arg Glu Gly Gly Val Cys Arg Thr Arg Thr Met Lys Ile	
125 130 135	
ATC ATG AAG GTT GGG CAA GAT CCC AAT GCT GTG ACG CCT GAG CAG CTG	528
Ile Met Lys Val Gly Gln Asp Pro Asn Ala Val Thr Pro Glu Gln Leu	
140 145 150	
ACT ACC AGC AGG CCC AGC AAG GAG GCA GAC AAC ACT GTC AAG ATG GCC	576
Thr Thr Ser Arg Pro Ser Lys Glu Ala Asp Asn Thr Val Lys Met Ala	
155 160 165	
ACA CAG GCC CCT GGT AGT CGG GGC TCC CTG GGT GAC TCT GAT GGC AAG	624
Thr Gln Ala Pro Gly Ser Arg Gly Ser Leu Gly Asp Ser Asp Gly Lys	
170 175 180	

30

```

CAT GAG ACT GTG AAC CAG GAA GAG AAG AGT GGC CCA GGT GCA AGT GGG 672
His Glu Thr Val Asn Gln Glu Glu Lys Ser Gly Pro Gly Ala Ser Gly
185                      190                      195                      200

GGC AGC AGC GGG GAC CCT GAT GGC TTC TTC AAC TCC AAG GTG GCA TTG 720
Gly Ser Ser Gly Asp Pro Asp Gly Phe Phe Asn Ser Lys Val Ala Leu
205                      210                      215

TTC GCG GCT GTC GGT GCC GGT TGC GTC ATC TTC CTG CTC ATC ATC ATC 768
Phe Ala Ala Val Gly Ala Gly Cys Val Ile Phe Leu Leu Ile Ile Ile
220                      225                      230

TTC CTG ACG GTC CTA CTA CTG AAG CTA CGC AAG CGG CAC CGC AAG CAC 816
Phe Leu Thr Val Leu Leu Leu Lys Leu Arg Lys Arg His Arg Lys His
235                      240                      245

ACA CAG CAG CGG GCG GCT GCC CTC TCG CTC AGT ACC CTG GCC AGT CCC 864
Thr Gln Gln Arg Ala Ala Ala Leu Ser Leu Ser Thr Leu Ala Ser Pro
250                      255                      260

AAG GGG GGC AGT GGC ACA GCG GGC ACC GAG CCC AGC GAC ATC ATC ATT 912
Lys Gly Gly Ser Gly Thr Ala Gly Thr Glu Pro Ser Asp Ile Ile Ile
265                      270                      275                      280

CCC TTA CGG ACT ACA GAG AAC AAC TAC TGC CCC CAC TAT GAG AAG GTG 960
Pro Leu Arg Thr Thr Glu Asn Asn Tyr Cys Pro His Tyr Glu Lys Val
285                      290                      295

AGT GGG GAC TAC GGG CAC CCT GTC TAC ATC GTC CAA GAG ATG CCG CCC 1008
Ser Gly Asp Tyr Gly His Pro Val Tyr Ile Val Gln Glu Met Pro Pro
300                      305                      310

CAG AGC CCG GCG AAC ATC TAC TAC AAG GTC TGA 1041
Gln Ser Pro Ala Asn Ile Tyr Tyr Lys Val
315                      320

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 346 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Ala Arg Pro Gly Gln Arg Trp Leu Gly Lys Trp Leu Val Ala Met
-24                      -20                      -15                      -10

Val Val Trp Ala Leu Cys Arg Leu Ala Thr Pro Leu Ala Lys Asn Leu
-5                      1                      5

Glu Pro Val Ser Trp Ser Ser Leu Asn Pro Lys Phe Leu Ser Gly Lys
10                      15                      20

Gly Leu Val Ile Tyr Pro Lys Ile Gly Asp Lys Leu Asp Ile Ile Cys
25                      30                      35                      40

```

Pro Arg Ala Glu Ala Gly Arg Pro Tyr Glu Tyr Tyr Lys Leu Tyr Leu
 45 50 55
 Val Arg Pro Glu Gln Ala Ala Ala Cys Ser Thr Val Leu Asp Pro Asn
 60 65 70
 Val Leu Val Thr Cys Asn Arg Pro Glu Gln Glu Ile Arg Phe Thr Ile
 75 80 85
 Lys Phe Gln Glu Phe Ser Pro Asn Tyr Met Gly Leu Glu Phe Lys Lys
 90 95 100
 His His Asp Tyr Tyr Ile Thr Ser Thr Ser Asn Gly Ser Leu Glu Gly
 105 110 115 120
 Leu Glu Asn Arg Glu Gly Gly Val Cys Arg Thr Arg Thr Met Lys Ile
 125 130 135
 Ile Met Lys Val Gly Gln Asp Pro Asn Ala Val Thr Pro Glu Gln Leu
 140 145 150
 Thr Thr Ser Arg Pro Ser Lys Glu Ala Asp Asn Thr Val Lys Met Ala
 155 160 165
 Thr Gln Ala Pro Gly Ser Arg Gly Ser Leu Gly Asp Ser Asp Gly Lys
 170 175 180
 His Glu Thr Val Asn Gln Glu Glu Lys Ser Gly Pro Gly Ala Ser Gly
 185 190 195 200
 Gly Ser Ser Gly Asp Pro Asp Gly Phe Phe Asn Ser Lys Val Ala Leu
 205 210 215
 Phe Ala Ala Val Gly Ala Gly Cys Val Ile Phe Leu Leu Ile Ile Ile
 220 225 230
 Phe Leu Thr Val Leu Leu Leu Lys Leu Arg Lys Arg His Arg Lys His
 235 240 245
 Thr Gln Gln Arg Ala Ala Ala Leu Ser Leu Ser Thr Leu Ala Ser Pro
 250 255 260
 Lys Gly Gly Ser Gly Thr Ala Gly Thr Glu Pro Ser Asp Ile Ile Ile
 265 270 275 280
 Pro Leu Arg Thr Thr Glu Asn Asn Tyr Cys Pro His Tyr Glu Lys Val
 285 290 295
 Ser Gly Asp Tyr Gly His Pro Val Tyr Ile Val Gln Glu Met Pro Pro
 300 305 310
 Gln Ser Pro Ala Asn Ile Tyr Tyr Lys Val
 315 320

CLAIMS

What is claimed is:

- 5 1. An isolated DNA encoding an elk-L protein capable of binding elk, wherein said DNA comprises a nucleotide sequence that is at least 80% identical to a sequence selected from the group consisting of nucleotides 1-1041, 73-1041, 1-711, and 73-711 of SEQ ID NO:1.
- 10 2. An isolated DNA according to claim 1, wherein said DNA comprises a nucleotide sequence selected from the group consisting of nucleotides 1-1041, 73-1041, 1-711, and 73-711 of SEQ ID NO:1.
- 15 3. An isolated DNA encoding an elk-L protein capable of binding elk, wherein said elk-L comprises an amino acid sequence that is at least 80% identical to a sequence selected from the group consisting of amino acids 1-322 and 1-213 of SEQ ID NO:2.
- 20 4. An isolated DNA according to claim 3, wherein said elk-L comprises an amino acid sequence selected from the group consisting of amino acids 1-322 and 1-213 of SEQ ID NO:2.
5. An expression vector comprising a DNA sequence according to claim 1, 2, 3, or 4.
- 25 6. A process for preparing an elk-L polypeptide, comprising culturing a host cell transformed with a vector according to claim 5 under conditions promoting expression of elk-L, and recovering the elk-L polypeptide from the culture.
- 30 7. Purified biologically active mature elk-L protein characterized by the N-terminal amino acid sequence Ala-Thr-Pro-Leu-Ala-Lys-Asn-Leu-Glu-Pro-Val-Ser-.
8. Purified elk-L according to claim 7, wherein said elk-L comprises an amino acid sequence selected from the group consisting of amino acids 1-322 and 1-213 of SEQ ID NO:2.

9. A purified elk-L protein comprising an amino acid sequence that is at least 80% identical to a sequence selected from the group consisting of amino acids 1-322 and 1-213 of SEQ ID NO:2.

5 10. An antibody that is immunoreactive with an elk-L protein according to claim 8, or with an immunogenic fragment thereof.

11. An antibody according to claim 10, wherein said antibody is a monoclonal antibody.

10

12. An isolated nucleic acid molecule comprising a sequence of at least about 14 nucleotides of the nucleotide sequence of SEQ ID NO:1 or the DNA or RNA complement thereof.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/10955**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) :C07H 21/04; C12N 15/12; C07K 13/00, 15/28

US CL :536/23.5, 24.31, 24.33, 24.5; 435/69.1; 530/350, 387.9, 388.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.5, 24.31, 24.33, 24.5; 435/69.1; 530/350, 387.9, 388.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE

search terms: elk, protein tyrosine kinase and ligand

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Molecular and Cellular Biology, Volume 11, No. 5, issued May 1991, V. Lhotak et al., "Characterization of Elk, a brain-specific receptor tyrosine kinase", pages 2496-2502, entire document.	1-12



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

03 February 1994

Date of mailing of the international search report

09 FEB 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

LORRAINE M. SPECTOR, PH.D.

Telephone No. (703) 308-0196

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.